

Generalization and Alternatives of Kaprekar's Routine

Florentin Smarandache
University of New Mexico, Gallup Campus, USA

Abstract.

We extend Kaprekar's Routine for a large class of applications. We also give particular examples of this generalization as alternatives to Kaprekar's Routine and Number. Some open questions about the length of the iterations until reaching either zero or a constant or a cycle, and about the length of the cycles are asked at the end.

1. Generalization of Kaprekar's Routine.

Let f be an operator that maps a finite set $A = \{a_1, a_2, \dots, a_p\}$, with $p \geq 1$, into itself:

$$f: A \rightarrow A.$$

Then, for any value $a \in A$ we have $f(a) \in A$ too. If we iterate this operator multiple times we get a chain like this:

$a \in A$ involves $f(a) \in A$, which involves $f(f(a)) \in A$ (f iterated with itself twice), which in its turn involves $f(f(f(a))) \in A$, and so on, $f(\dots f(a) \dots) \in A$ { where, in the last case, f has been iterated with itself $r \geq 1$ times; let's denote it by $f_r(a)$ }. Let's also denote, for the symmetry of notation,

$$a = f_0(a).$$

Since cardinal of A is a finite positive integer, $\text{card}(A) = p < +\infty$, after at most $r = p$ iterations we get two equal iterations

$$f_i(a) = f_j(a)$$

for $i \neq j$ with $0 \leq i < j \leq r$, in the above chain $a = f_0(a), f_1(a), f_2(a), \dots, f_r(a)$.

Hence, this chain of values can form a cycle:

$$f_0(a), f_1(a), \dots, f_i(a), f_{i+1}(a), \dots, f_{j-1}(a), f_i(a), f_{i+1}(a), \dots, f_{j-1}(a),$$

with the cycle $f_i(a), f_{i+1}(a), \dots, f_{j-1}(a)$ of length $j-i$.

If $j=i+1$ then the chain reaches a constant, since the cycle has only one element $f_i(a)$.

2. Kaprekar's Routine.

Kaprekar's Routine {see [1], [2], [3]}, extended to k -digit numbers, is a particular case of the above algorithm.

For $k = 1$ and 2 the Kaprekar's Routine reached zero, for $k = 3$ or 4 the algorithm reaches zero or 495, respectively zero or 6174. For $k \geq 6$ it also reaches cycles.

The set $A = \{0, \text{ and all } k\text{-digit positive integers}\}$, so its cardinal is finite, $\text{card}(A) = 90 \dots 01$, where in this number we have $k-2$ zeroes.

The operator f does the following: arranges the digits of number a in descending order (a') and in ascending order (a'') and then subtracts them: $a' - a''$. Since $a' - a''$ is also a k -digit number or zero (in the degenerate case when all the digits are equal), then $f(a' - a'')$ is a k -digit number as well or zero, therefore $f(a' - a'') \in A$. And the iteration continues in the same way. After a finite number of iterations the algorithm reaches a constant (which can be zero in the degenerate case), or a constant, or gets into a cycle.

3. Alternatives to Kaprekar's Routine.

3.1. Let's consider the group of permutations P of the digits of the k -digit ($k \geq 1$) number a .

We define the operator

$$f_P(a) = |P_1(a) - P_2(a)|$$

where P_1 and P_2 are some permutations of k elements $\{1, 2, \dots, k\}$, and $|\cdot|$ means absolute value.

And the set $A = \{0; \text{ and all } m\text{-digit positive integers, } m \leq k\}$.

Then the sequence of iterations reaches zero, a constant, or a cycle.

Let's see an **example**:

$P_1(\{1, 2, 3\}) = \{2, 3, 1\}$ and $P_2(\{1, 2, 3\}) = \{1, 3, 2\}$; $a = 125$, then we have $|251 - 152| = 099$;

$|990 - 099| = 891$; $|918 - 819| = 099$; $|990 - 099| = 891$; $|918 - 819| = 099$;

So, we reached a cycle: 125, 099, 891, 099, 891,

3.2. Let's have the same group of permutations and same set A as in Example 3.1, but taking as operator:

$$fa_P(a) = |a - P(a)|.$$

See another example:

$P(\{1,2,3\}) = \{3,1,2\}$; $a = 125$, then $|125-512| = 387$, $|387-738| = 351$, $|351-135| = 216$,
 $|216-621| = 405$, $|405-540| = 135$, $|135-513| = 378$, $|378-837| = 459$, $|459-945| = 486$,
 $|486-648| = 162$, $|162-216| = 054$, $|054-405| = 351$,

So, we got: 125, 387, 351, 216, 405, 135, 378, 459, 486, 162, 054, 351,

3.3. Similarly if we take the operator: the absolute value of a number minus its reverse:

$$|a - \text{reverse}(a)|.$$

For example: 125, $|125-521| = 396$, $|396-693| = 297$, $|297-792| = \underline{495}$, $|495-594| = 099$,
 $|099-990| = 891$, $|891-198| = 693$, $|693-396| = 297$, $|297-792| = \underline{495}$,

3.4. Let's consider the *Smarandache Form* of a number. A *Smarandache Palindrome* (SP) {see [4]-[13]} is a number of the form $a_1a_2...a_{n-1}a_na_na_{n-1}...a_2a_1$ or $a_1a_2...a_{n-1}a_na_{n-1}...a_2a_1$ where each a_i is a positive integer of any number of digits. For example, 143431 is a SP since it can be written as 1(43)(43)1. A Smarandache Form is any number $a_1a_2...a_{n-1}a_n$, where each a_i is a positive integer of any number of digits.

Then we can take the operator mapping SFs into SFs in the following way.

Example: Consider the following SF: 3-digit numbers under the SF of 1-digit and 2-digit groups: for example 5(76). Then we switch the groups and add them. Take modulo 1000 of the result.

Start with 5(76), $5(76) + (76)5 = 1342$ whole modulo 1000 is $342 = 3(42)$;
 $3(42) + (42)3 = 767 = 7(67)$; $7(67) + (67)7 = 1444$ whose modulo 1000 is $444 = 4(44)$;
 $4(44) + (44)4 = 888 = 8(88)$; $8(88) + (88)8 = 1776$ whose modulo 1000 is $776 = 7(76)$;
 $7(76) + (76)7 = 1543$ whose modulo 1000 is $543 = 5(43)$;
 $5(43) + (43)5 = 978$; $9(78) + (78)9 = 1767$ whose modulo 1000 is $767 = 7(67)$;
 $7(67) + (67)7 = 1444$ whose modulo 1000 is $4(44)$.

We got: 5(76), 3(42), 4(44), 7(76), 5(43), 7(67), 4(44),

3.5. Or one consider another operator that subtracts two k -digit numbers in the following way: adding 1 to each digit less than 9, then subtracting 1 from each non-zero digit, then subtracting the numbers.

Example: $495, 596-384 = 212, 323 - 101 = \underline{222}, 333-111 = \underline{222}, \dots$. We reached a constant.

Or add 2 to each digit strictly less than 8, and subtract 3 from each digit strictly greater than 2. Etc.

3.6. Consider any function f defined on the set of k -digit numbers whose range is the set of positive integers, and then calculate *modulo* 10^k of the result.

There are infinitely many such operators in order to choose from nice examples.

4. Open Questions:

1. What is the longest number of iterations until one reaches either zero or a constant or a cycle that one can have for each case of the above generalizations?
2. What is the longest cycle that one can have for each particular case of the above generalizations?
3. In what conditions one reaches a constant, not a cycle? By cycle we understand a sequence of two or more numbers that repeat indefinitely.
4. Study the cases when $f(a) = a$ for interesting particular cases of this generalization.
5. Study the case when $f(a) = 0$ for interesting particular cases of this generalization.
6. If the operator defined in the above Generalization of the Kaprekar's Routine is a random operator (i.e. for a given k -digit number a one randomly generate another k -digit number b), is it still possible to reach a constant or a cycle?

It is possible for sure to generate two equal k -digit numbers, c , after at most 10^k random operations, but then the next k -digit number following the first c would not necessarily be the same as the previous k -digit number following the previous c .

References:

1. D. R. Kaprekar, *An Interesting Property of the Number 6174*, Scripta Math., Vol. 15, 244-245, 1955.
2. N. J. A. Sloane, *Sequence A099009*, Online Encyclopedia of Integers, <http://www.research.att.com/~njas/sequences/A099009>.
3. Eric W. Weisstein, *Kaprekar Routine*, MathWorld – A Wolfram Web Resource, <http://mathworld.wolfram.com/KaprekarRoutine.html>.
4. Florentin Smarandache, *Properties of Numbers*, Arhivele Statului, Filiala Vâlcea, Rm. Vâlcea, 1972.
5. Charles Ashbacher, Lori Neiryneck, *The Density of Generalized Smarandache Palindromes*, Journal of Recreational Mathematics, Vol. 33 (2), 2006.
6. G. Gregory, *Generalized Smarandache Palindromes*, <http://fs.gallup.unm.edu//GSP.htm>.
7. M. Khoshnevisan, "Generalized Smarandache Palindrome", Mathematics Magazine, Aurora, Canada, 10/2003.
8. N. Sloane, *Encyclopedia of Integers*, Sequence A082461, <http://www.research.att.com/cgi-bin/access.cgi/as/njas/sequences/eisA.cgi?Anum=A082461>
9. F. Smarandache, *Sequences of Numbers Involved in Unsolved Problems*, Hexis, 1990, 2006; <http://www.gallup.unm.edu/~smarandache/Sequences-book.pdf>.

10. Temitope Gbolahan Jaiyeola, *Palindromic permutations and generalized Smarandache palindromic permutations*, Scientia Magna, 2006.
11. Planet Math,
<http://planetmath.org/encyclopedia/GeneralizedSmarandachePalindrome.html>.
12. M. Khoshnevisan, Proposed Problem #1062 (on *Generalized Smarandache Palindrome*), The ΠME Epsilon, USA, Vol. 11, No. 9, p. 501, Fall 2003.
13. Mark Evans, Mike Pinter, Carl Libis, Solutions to Problem #1062 (on *Generalized Smarandache Palindrome*), The ΠME Epsilon, Vol. 12, No. 1, 54-55, Fall 2004.

{Partially published in author's book *Proposed Problems of Mathematics, Vol. II*, State University of Moldova Press, Kishinev, pp. 83-84, 1997.}